

# Lecture 13: The Curry-Howard-Lambek correspondence for classical computation

## Summary.

- (1) The Curry-Howard-Lambek correspondence: from logic to categories and back.
- (2) The Curry-Howard-Lambek correspondence: from programs to categories and back.


*Lúis Soares Barbosa,*

UNIV. MINHO (*Informatics Dep.*) & INL (*Quantum Software Engineering Group*)


## Overview.

---

The general triangle



is instantiated to



A previous lecture already discussed the link between (intuitionistic) logic and (simply-typed)  $\lambda$ -calculus under the *motto*

Formulas-as-Types and Proofs-as-Programs

It was emphasized that exploring the computational content of proofs is, indeed, fully aligned with the constructive (BHK) interpretation of intuitionistic logic under which, for example, a proof of  $A \wedge B$  is a pair of proofs of both  $A$  and  $B$ , and a proof of  $A \longrightarrow B$  is a procedure to transform any proof of  $A$  into a proof of  $B$ . We turn now to the links that both logic and computation keep with the mathematical structures which provide their semantical models, i.e with *categories*.

Given a Cartesian-closed category (CCC)  $\mathcal{C}$ , the Lambek's part of the diagram identifies

Formulas-as-Objects and Proofs-as-Arrows

Recall the basic structure of a CCC:

- *Products*:  $A \times B$ , with projections  $\pi_1, \pi_2$  and a split arrow  $\langle f, g \rangle : C \rightarrow A \times B$  defined by a universal property from  $f : C \rightarrow A$  and  $g : C \rightarrow B$ . The product construction is functorial:  $f \times g = \langle f \cdot \pi_1, g \cdot \pi_2 \rangle$ .
- *Exponentials*:  $B^A$ , given through the natural isomorphism between

$$f : A \times B \rightarrow C \quad \Leftrightarrow \quad \bar{f} : A \rightarrow C^B$$

expressed through another universal property

$$k = \bar{f} \quad \Leftrightarrow \quad f = \text{ev} \cdot (k \times \text{id})$$

$$\begin{array}{ccc}
 A \times B & \xrightarrow{\bar{f} \times \text{id}} & C^B \times B \\
 & \searrow f & \downarrow \text{ev} \\
 & & Y
 \end{array}$$

$$A \xrightarrow{\bar{f}} C^B$$

Construction  $-^C$  extends to a functor, the covariant *exponential* functor, by defining

$$h^C : X^C \rightarrow Y^C = \overline{(h \cdot \text{ev})}$$

for  $h : X \rightarrow Y$ .

Note that the exponential object  $X^C$  represents as an object in the category, the arrows from  $C$  to  $X$ . Consequently, the action of  $-^C$  on each arrow  $f : X \rightarrow Y$  should *internalise* composition with  $h$ . In *Set* it is easy to verify that this is indeed the case. For  $g : C \rightarrow X$  and  $c \in C$ , a simple calculation yields,

$$\begin{aligned}
 & h^C(g)(c) \\
 = & \{ h^C = \overline{(h \cdot \text{ev})} \} \\
 & \overline{(h \cdot \text{ev})(g)}(c) \\
 = & \{ \text{uncurrying} \} \\
 & h \cdot \text{ev}(g, c) \\
 = & \{ \text{function composition} \}
 \end{aligned}$$

$$\begin{aligned}
& \mathbf{h}(\mathbf{ev}(g, c)) \\
= & \quad \{ \text{ev definition} \} \\
& \mathbf{h}(g(c)) \\
= & \quad \{ \text{function composition} \} \\
& (\mathbf{h} \cdot g)(c)
\end{aligned}$$

which means that  $\mathbf{h}^{\mathcal{C}} = \mathbf{h} \cdot \dots$

---

### The link to logic.

---

Formulas in intuitionistic logic correspond to objects in  $\mathcal{C}$ ; proofs correspond to morphisms in  $\mathcal{C}$ . The correspondence is as follows:

Intuitionistic logic	CCC
$\frac{}{\Gamma, x : A \vdash A}$	$\frac{}{\pi_2 : \Gamma \times A \longrightarrow A}$
$\frac{\Gamma \vdash A \quad \Gamma \vdash B}{\Gamma \vdash A \wedge B}$	$\frac{f : \Gamma \longrightarrow A \quad g : \Gamma \longrightarrow B}{\langle f, g \rangle : \Gamma \longrightarrow A \times B}$
$\frac{\Gamma \vdash A \wedge B}{\Gamma \vdash A}$	$\frac{f : \Gamma \longrightarrow A \times B}{\pi_1 \cdot f : \Gamma \longrightarrow A}$
$\frac{\Gamma \vdash A \wedge B}{\Gamma \vdash B}$	$\frac{f : \Gamma \longrightarrow A \times B}{\pi_2 \cdot f : \Gamma \longrightarrow B}$
$\frac{\Gamma, A \vdash B}{\Gamma \vdash A \longrightarrow B}$	$\frac{f : \Gamma \times A \longrightarrow B}{\bar{f} : \Gamma \longrightarrow B^A}$
$\frac{\Gamma \vdash A \longrightarrow B \quad \Gamma \vdash A}{\Gamma \vdash B}$	$\frac{f : \Gamma \longrightarrow B^A \quad g : \Gamma \longrightarrow A}{\mathbf{ev}_{A,B} \cdot \langle f, g \rangle : \Gamma \longrightarrow B}$

#### Exercise 1

Extend the CHL correspondence to capture the propositional intuitionistic logic is enriched with disjunction, i.e. connectives  $\vee$  and  $\perp$ .

---

## The link to computation.

---

Types-as-Objects and Programs-as-Arrows

Types in the simply-typed  $\lambda$ -calculus correspond objects in a CCC  $\mathcal{C}$ . Programs, i.e. terms in the simply-typed  $\lambda$ -calculus, on the other hand, correspond to morphisms in  $\mathcal{C}$ . Moreover, the  $\beta, \eta$ -reduction is suitably derived from the axioms of a CCC. The correspondence is captured by a semantic function which translates each term

$$x_1 : A_1, \dots, x_n : A_n \vdash u : B$$

into an arrow in  $\mathcal{C}$ :

$$\llbracket u \rrbracket : \llbracket A_1 \rrbracket \times \dots \times \llbracket A_n \rrbracket \longrightarrow \llbracket B \rrbracket$$

The correspondence is defined recursively on types by

$$\begin{aligned} \llbracket A \times B \rrbracket &\hat{=} \llbracket A \rrbracket \times \llbracket B \rrbracket \\ \llbracket A \longrightarrow B \rrbracket &\hat{=} \llbracket B \rrbracket^{\llbracket A \rrbracket} \end{aligned}$$

assuming a set of distinguished objects in  $\mathcal{C}$  as semantic domains for the basic types.

Similarly, for terms,

$$\frac{}{\llbracket \Gamma, x : A \vdash x : A \rrbracket \hat{=} \pi_2 : \llbracket \Gamma \rrbracket \times \llbracket A \rrbracket \longrightarrow \llbracket A \rrbracket}$$

$$\llbracket \Gamma \vdash u : A \times B \rrbracket = f : \llbracket \Gamma \rrbracket \longrightarrow \llbracket A \rrbracket \times \llbracket B \rrbracket$$

$$\frac{}{\llbracket \Gamma \vdash \pi_1 u : A \rrbracket \hat{=} \pi_1 \cdot f : \llbracket \Gamma \rrbracket \longrightarrow \llbracket A \rrbracket}$$

$$\frac{\llbracket \Gamma \vdash u : A \rrbracket = f : \llbracket \Gamma \rrbracket \longrightarrow \llbracket A \rrbracket \quad \llbracket \Gamma \vdash v : B \rrbracket = g : \llbracket \Gamma \rrbracket \longrightarrow \llbracket B \rrbracket}{\llbracket \Gamma \vdash \langle u, v \rangle : A \times B \rrbracket \hat{=} \langle f, g \rangle : \llbracket \Gamma \rrbracket \longrightarrow \llbracket A \rrbracket \times \llbracket B \rrbracket}$$

$$\frac{}{\llbracket \Gamma, x : A \vdash u : B \rrbracket = f : \llbracket \Gamma \rrbracket \times \llbracket A \rrbracket \longrightarrow \llbracket B \rrbracket}$$

$$\frac{}{\llbracket \Gamma \vdash \lambda x. u : A \longrightarrow B \rrbracket \hat{=} \bar{f} : \llbracket \Gamma \rrbracket \longrightarrow \llbracket B \rrbracket^{\llbracket A \rrbracket}}$$

$$\frac{\llbracket \Gamma \vdash u : A \longrightarrow B \rrbracket = f \quad \llbracket \Gamma \vdash v : A \rrbracket = g}{\llbracket \Gamma \vdash uv : B \rrbracket \hat{=} ev \cdot \langle f, g \rangle : \llbracket A \rrbracket \longrightarrow \llbracket B \rrbracket}$$

---

## Soundness of $\llbracket - \rrbracket$ .

Soundness of the translation of simply-typed  $\lambda$ -calculus to a CCC means that  $\beta, \eta$ -equivalence, which equates terms that are derived one from the other through the rules of  $\beta, \eta$ -reduction, correspond to semantic equality, i.e.

$$\boxed{u =_{\beta, \eta} v \Rightarrow \llbracket u \rrbracket = \llbracket v \rrbracket}$$

Let  $\Gamma = x_1 : A_1 \cdots A_n$ . Given terms  $\Gamma \vdash u : A$  and, for all  $1 \leq i \leq n$ ,  $\Gamma \vdash u_i : A_i$ ,

$$\llbracket u[x_1 := u_1, \dots, x_n := u_n] \rrbracket = \llbracket u \rrbracket \cdot \langle \llbracket u_1 \rrbracket \cdots \llbracket u_n \rrbracket \rangle$$

This statement, known as the *substitution lemma*, is proved by induction on the structure of terms. The base case is that of variables:  $x_i$ . Actually,

$$\llbracket x_i[x := u] \rrbracket = \llbracket u_i \rrbracket = \pi_i \cdot \langle \llbracket u_1 \rrbracket, \dots, \llbracket u_k \rrbracket \rangle = \llbracket x_i \rrbracket \cdot \langle \llbracket u_1 \rrbracket, \dots, \llbracket u_k \rrbracket \rangle$$

For the inductive process, consider, for example,  $\lambda x. u$ . Thus,

$$\begin{aligned} & \llbracket \lambda x. u[x := v] \rrbracket \\ = & \{ \text{substitution} \} \\ & \llbracket \lambda x. u[x, x := v, x] \rrbracket \\ = & \{ \llbracket - \rrbracket \text{ definition} \} \\ & \overline{\llbracket u[x, x := v, x] \rrbracket} \\ = & \{ \text{induction hypothesis} \} \\ & \overline{\llbracket u \rrbracket \cdot (\langle v \rangle \times \text{id})} \\ = & \{ \text{fusion law for exponentials: } \bar{f} \cdot g = \overline{f \cdot (g \times \text{id})} \} \\ & \overline{\llbracket u \rrbracket} \cdot \langle v \rangle \\ = & \{ \llbracket - \rrbracket \text{ definition} \} \\ & \llbracket \lambda x. u \rrbracket \cdot \langle v \rangle \end{aligned}$$

**Exercise 2**

Complete the proof of the *substitution lemma* above for the remaining cases.

---

To establish soundness of the semantic interpretation  $\llbracket \_ \rrbracket$ , all we need to show is that the interpretation of both sides of a  $\beta, \eta$ -reduction corresponds to a valid equation in a CCC. The substitution lemma is an important tool in this proof.

Let us start with  $\beta$ -conversion, considering the interpretation of

$$\begin{aligned}
 (\lambda x . u)v &=_{\beta} u[x := v] \\
 \llbracket (\lambda x . u)v \rrbracket &= \{ \llbracket \_ \rrbracket \text{ definition} \} \\
 &= ev \cdot \langle \overline{\llbracket u \rrbracket}, \llbracket v \rrbracket \rangle \\
 &= \{ \times\text{-absorption law} \} \\
 &= ev \cdot (\overline{\llbracket u \rrbracket} \times id) \cdot \langle id, \llbracket v \rrbracket \rangle \\
 &= \{ \text{currying definition} \} \\
 &= \llbracket u \rrbracket \cdot \langle id, \llbracket v \rrbracket \rangle \\
 &= \{ \text{substitution lemma} \} \\
 &= \llbracket u[x, v := x, x] \rrbracket
 \end{aligned}$$

**Exercise 3**

Verify the second  $=_{\beta}$ -conversion

$$\pi_1 \langle u, v \rangle = u \quad \text{and} \quad \pi_2 \langle u, v \rangle = v$$


---

**Exercise 4**

Verify the two  $=_{\eta}$ -conversions

$$u = \lambda x . u x$$

and

$$u = \langle \pi_1 u, \pi_2 u \rangle$$


---

## Completeness of $\llbracket - \rrbracket$ .

To show completeness one has to come up with a concrete CCC,  $\Lambda$ , in which equalities between arrows correspond to  $\beta, \eta$ -conversions between terms, i.e.

$$\boxed{u =_{\beta, \eta} v \iff \llbracket u \rrbracket = \llbracket v \rrbracket}$$

where  $\llbracket - \rrbracket$  is an interpretation of  $\lambda$ -terms in  $\Lambda$ .

The category  $\Lambda$  has an object  $\hat{A}$  for each type  $A$  in the  $\lambda$ -calculus, plus a final object  $\mathbf{1}$ . An arrow from  $\hat{A}$  to  $\hat{B}$  is an equivalence class of the following relation defined on variable-term pairs:

$$(x, u) \approx (y, v) \quad \text{iff} \quad x : A \vdash u : B \quad \text{and} \quad y : A \vdash v : B \quad \text{and} \quad u =_{\beta, \eta} v[y := x]$$

which extends to pairs  $(*, u)$ , where  $*$  represents the single inhabitant of  $\mathbf{1}$ , as follows:

$$(*, u) \approx (*, v) \quad \text{iff} \quad \vdash u : B \quad \text{and} \quad \vdash v : B \quad \text{and} \quad u =_{\beta, \eta} v$$

As usual, the equivalence class  $\llbracket (x, u) \rrbracket$ , for the element  $(x, u)$ , is the set  $\{(y, v) \mid (x, u) \approx (y, v)\}$ . Thus, the homsets of  $\Lambda$  are as follows:

$$\begin{aligned} \Lambda[\hat{A}, \hat{B}] &= \{\llbracket (x, u) \rrbracket \mid x : A \vdash u : B\} \\ \Lambda[\mathbf{1}, \hat{B}] &= \{\llbracket (*, u) \rrbracket \mid \vdash u : B\} \\ \Lambda[\hat{A}, \mathbf{1}] &= \{!_{\hat{A}}\} \end{aligned}$$

### Exercise 5

In  $\Lambda$  define,

- Identities:  $\text{id}_{\hat{A}} \hat{=} \llbracket (x, x) \rrbracket$  and  $\text{id}_{\mathbf{1}} \hat{=} !_{\mathbf{1}}$
- Composition:

$$\begin{aligned} \llbracket (x, u) \rrbracket \cdot \llbracket (y, v) \rrbracket &\hat{=} \llbracket (y, u[x := v]) \rrbracket \\ \llbracket (x, u) \rrbracket \cdot \llbracket (*, v) \rrbracket &\hat{=} \llbracket (*, u[x := v]) \rrbracket \\ \llbracket (*, u) \rrbracket \cdot !_Z &\hat{=} \begin{cases} \llbracket (y, u) \rrbracket & \Leftarrow Z = \hat{A} \\ \llbracket (*, u) \rrbracket & \Leftarrow Z = \mathbf{1} \end{cases} \\ !_W \cdot h &\hat{=} !_Z \quad \text{for } h : Z \longrightarrow W \end{aligned}$$

Prove that  $\Lambda$  is a category.

The category  $\Lambda$  has finite products and exponentials, and provides what is called a *term* (i.e. built on top of the syntax) model for the simply-typed  $\lambda$ -calculus (see, e.g. [1] for proofs).

## References

- [1] S. Abramsky and N. Tzevelekos. Introduction to categories and categorical logic. In B. Coecke, editor, *New Structures for Physics*, pages 3–94. Springer Lecture Notes on Physics (813), 2011.